



IPM PARTNER®

Please Distribute

Field Reporter™

Your Edge --- And Ours ---- Is Knowledge

Volume 2

IPM PARTNER® FIELD REPORTER March 4, 2010

NOTE: Oriental Fruit Moth (OFM) – Moth catch is expected in all areas as soon as weather warms. Traps should be in place for that warm weather.

Kern County (Delano/McFarland): **OFM** - Some moths trapped 3/1.

Tulare County (NE Kingsburg area): **OFM** - No activity

Madera County: **OFM** - No activity

Merced County (Ballico & Winton areas): **OFM** - A few moths trapped but no sustained flight

Sacramento County (Delta area): **OFM** - No activity

Yuba County (Marysville): **OFM** - No activity

Butte County (Gridley): **OFM** - No activity

NOTE: Mark your calendars for the dates listed below.

FEBRUARY: OFM, OLR and SJS traps should be set.

MARCH: CM, CM-DA COMBO, DA, CM 10X, CRS, DBM AND PTB traps should be set.

Trece is pleased to announce their latest scientific innovation, CIDETRAK® Mating Disruptant, an intelligently simple high performance dispenser for Codling Moth and Oriental Fruit Moth mating disruption. Also, don't forget PHEROCON® CM-DA COMBO multi-gender attractant for monitoring which will give you a definite edge in your battle against the codling moth, your toughest competition. This lure has been proven to be the most effective codling moth lure on the market today. For more information on CIDETRAK or DA and CM-DA COMBO lures, and how to interpret data from trap capture, contact Jeff Downs, National Sales Manager, at 559-285-8448.

The IPM PARTNER® Field Reporter™ will also be published on our website www.trece.com

	Cellular Phone	Office	E-Mail
Jeff Downs, National Sales Manager	(559) 285-8448	(480) 639-9495	jdowns@trece.com
Donna Lingren, Marketing Manager donna lingren@earthlink.net	(918) 261-1377	(918) 785-3061	
Debi Bilby, Customer Service Representative		(918) 785-3061	dbilby@trece.com

NOTE: The foregoing is supplied as a general information guideline only. This information is specific to the site location from which it was obtained. For optimum pest management decision making, each orchard site should be individually monitored with traps and environmental monitoring equipment.